



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Missed Anterior Inferior Cerebellar Artery Aneurysm Mimicking Vestibular Neuritis-Clues to Prevent Misdiagnosis

Willms, J F ; Baltsavias, G ; Burkhardt, J K ; Ernst, S ; Tarnutzer, A A

Abstract: We discuss a case with combined vestibulocochlear and facial neuropathy mimicking a less urgent peripheral vestibular pattern of acute vestibular syndrome (AVS). With initial magnetic resonance imaging read as normal, the patient was treated for vestibular neuropathy until headaches worsened and a diagnosis of subarachnoid hemorrhage was made. On conventional angiography, a ruptured distal right-sided aneurysm of the anterior inferior cerebellar artery was diagnosed and coiled. Whereas acute vestibular loss usually points to a benign peripheral cause of AVS, combined neuropathy of the vestibulocochlear and the facial nerve requires immediate neuroimaging focusing on the cerebellopontine angle. Imaging should be assessed jointly by neuroradiologists and the clinicians in charge to take the clinical context into account.

DOI: <https://doi.org/10.1016/j.jstrokecerebrovasdis.2016.09.027>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-127672>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Willms, J F; Baltsavias, G; Burkhardt, J K; Ernst, S; Tarnutzer, A A (2016). Missed Anterior Inferior Cerebellar Artery Aneurysm Mimicking Vestibular Neuritis-Clues to Prevent Misdiagnosis. *Journal of Stroke and Cerebrovascular Diseases*, 25(12):e231-e232.

DOI: <https://doi.org/10.1016/j.jstrokecerebrovasdis.2016.09.027>

Missed AICA aneurysm mimicking vestibular neuritis – clues to prevent misdiagnosis

Jan-Folkard Willms M.D.¹, Gerasimos Baltasavias M.D.², Jan-Karl Burkhardt M.D.¹, Silvia Ernst M.D.³, Alexander A. Tarnutzer M.D.¹

1 Department of Neurosurgery, University Hospital Zurich and University of Zurich, Frauenklinikstr. 10, 8091 Zurich, Switzerland

2 Department of Neuroradiology, University Hospital Zurich, Frauenklinikstr. 10, 8091 Zurich, Switzerland

3 Department of Internal Medicine, Hospital Uster, Brunnenstrasse 42, 8610 Uster, Switzerland

Running title: AICA aneurysm mimicking vestibular neuritis

Corresponding author:

Alexander A. Tarnutzer, MD. Department of Neurosurgery, University Hospital Zurich. Frauenklinikstr. 10, 8091 Zurich, Switzerland. Phone: 0041 44 255 11 11. Fax: 0041 44 255 43 80. Email: alexander.tarnutzer@usz.ch

Conflict of interest and source of funding: Dr. Willms, Dr. Baltasavias, Dr. Ernst, Dr. Burkhardt and Dr. Tarnutzer report no conflict of interest. This study did not receive any funding.

Submission date: September 12th 2016

Abstract

We discuss a case with combined vestibulo-cochlear and facial neuropathy mimicking a less urgent peripheral-vestibular pattern of acute vestibular syndrome (AVS). With initial MR-imaging read as normal, the patient was treated for vestibular neuropathy until headaches worsened and a diagnosis of subarachnoid hemorrhage was made. On conventional angiography a ruptured distal right-sided aneurysm of the anterior inferior cerebellar artery was diagnosed and coiled. While acute vestibular loss usually points to a benign peripheral cause of AVS, combined neuropathy of the vestibulo-cochlear and the facial nerve requires immediate neuroimaging focusing on the cerebellopontine angle. Imaging should be assessed jointly by neuroradiologists and the clinicians in charge to take the clinical context into account.

Key words: MRI; acute vestibular syndrome; aneurysm; anterior inferior cerebellar artery; and diagnostic errors

Manuscript

Acute dizziness/vertigo accounts for 3.3-4.4% of all emergency-department consultations.[1] If dizziness/vertigo persists >24h and is accompanied by nausea/vomitus, nystagmus and gait imbalance, it is called acute vestibular syndrome (AVS).[2] A bedside ocular-motor examination (H.I.N.T.S.-plus; Head-Impulse-test, Nystagmus, Test-of-Skew, hearing-loss) proved successful in identifying those 25±15% with central causes.[2, 3] Here we report on pitfalls in peripheral-type AVS.

An 81-year-old woman presented with progressive vertigo over days, nausea/vomitus, new-onset headache and gait imbalance. A spontaneous left-beating nystagmus and a right-sided peripheral facial palsy were noted. Head-impulse testing was inconclusive, hearing was not assessed. She was hospitalized for suspected stroke and received a contrast-enhanced MRI, which was read as normal. Vestibular neuritis was considered and prednisone was prescribed. The patient's condition stabilized. However, on day five her headache suddenly worsened and her GCS dropped to 7. Head-CT demonstrated acute subarachnoid hemorrhage (SAH) (Fig. 1A). On digital-subtraction angiography a ruptured aneurysm of the right anterior-inferior cerebellar artery (AICA) was identified (Fig. 1B). Retrospectively, the aneurysm could be recognized on the initial MR-images (Fig. 1CD), demonstrating close proximity to the vestibulo-cochlear nerve and the facial nerve. Endovascular treatment was performed (Fig. 1E). While neurologically stable, she soon developed abdominal pain. On emergency surgery mesenteric ischemia was found. Facing extensive necrosis, palliative treatment was chosen. She died the next day.

New-onset headache, facial palsy and the progressive course over several days are the essential clues. This combination requires early vigilance and immediate evaluation. Except for Ramsey-Hunt-syndrome (with characteristic vesicles behind the ear, within the auditory canal and the palate),[4] a peripheral-type facial palsy or hearing-loss are incompatible with vestibular neuritis and must be considered 'red-flags'. [3] With the H.I.N.T.S.-plus likely

being negative and clinical findings pointing to the cerebello-pontine angle, MR-imaging and joint-assessment by the neuroradiologist and the clinician is paramount. While bleeding into a vestibular schwannoma may be considered,[5] dangerous vascular causes must be excluded.

Incidence for intracranial aneurysms is 9.7-14.5/100'000,[6] resulting in ~30'000 SAH annually in the US. AICA-aneurysms represent a tiny fraction (1-2%),[7] with distal location being exceptional (0.03%-0.22%, ~30 cases annually).[7, 8] While ~80% are diagnosed after rupture,[9] cerebello-pontine signs may be observed in unruptured AICA-aneurysms.[7, 9] This includes sudden hearing-loss or vertigo followed by facial palsy,[10] isolated acute,[8] episodic or chronic vertigo,[11] misdiagnosed as vestibular schwannoma or other cerebello-pontine-angle tumors.[10, 11] Warning signs in distal AICA-aneurysms may be observed in up to 50%.[8] With 250'000-500'000 AVS-cases annually,[2] distal AICA-aneurysms reflect <0.01%, making it a rare, but dangerous differential diagnosis. There should be a high-index of suspicion for vascular events, as current imaging techniques are limited for the inner ear and such changes are easily missed, sometimes requiring repeated imaging.

Required statements

The next of kin (son of the deceased patient) has consented to submission of this case report to the journal.

References

- [1] Newman-Toker DE, Hsieh YH, Camargo CA, Jr., Pelletier AJ, Butchy GT, Edlow JA. Spectrum of dizziness visits to US emergency departments: cross-sectional analysis from a nationally representative sample. *Mayo Clin Proc.* 2008;83:765-75.
- [2] Tarnutzer AA, Berkowitz AL, Robinson KA, Hsieh YH, Newman-Toker DE. Does my dizzy patient have a stroke? A systematic review of bedside diagnosis in acute vestibular syndrome. *CMAJ.* 2011;183:E571-92.
- [3] Newman-Toker DE, Kerber KA, Hsieh YH, Pula JH, Omron R, Saber Tehrani AS, et al. HINTS outperforms ABCD2 to screen for stroke in acute continuous vertigo and dizziness. *Acad Emerg Med.* 2013;20:986-96.
- [4] Iwasaki H, Toda N, Takahashi M, Azuma T, Nakamura K, Takao S, et al. Vestibular and cochlear neuritis in patients with Ramsay Hunt syndrome: a Gd-enhanced MRI study. *Acta Otolaryngol.* 2013;133:373-7.
- [5] Niknafs YS, Wang AC, Than KD, Etame AB, Thompson BG, Sullivan SE. Hemorrhagic vestibular schwannoma: review of the literature. *World Neurosurg.* 2014;82:751-6.
- [6] Connolly ES, Jr., Rabinstein AA, Carhuapoma JR, Derdeyn CP, Dion J, Higashida RT, et al. Guidelines for the management of aneurysmal subarachnoid hemorrhage: a guideline for healthcare professionals from the American Heart Association/american Stroke Association. *Stroke.* 2012;43:1711-37.
- [7] Gonzalez LF, Alexander MJ, McDougall CG, Spetzler RF. Anteroinferior cerebellar artery aneurysms: surgical approaches and outcomes--a review of 34 cases. *Neurosurgery.* 2004;55:1025-35.
- [8] Tokimura H, Ishigami T, Yamahata H, Yonezawa H, Yokoyama S, Haruzono A, et al. Clinical presentation and treatment of distal anterior inferior cerebellar artery aneurysms. *Neurosurg Rev.* 2012;35:497-503.
- [9] Yamakawa H, Hattori T, Tanigawara T, Sahashi Y, Ohkuma A. Intracanalicular aneurysm at the meatal loop of the distal anterior inferior cerebellar artery: a case report and review of the literature. *Surg Neurol.* 2004;61:82-8.
- [10] Diaz RC, Konia T, Brunberg J. Labyrinthine artery aneurysm as an internal auditory canal mass. *J Neurol Surg Rep.* 2014;75:e38-41.
- [11] Zager EL, Shaver EG, Hurst RW, Flamm ES. Distal anterior inferior cerebellar artery aneurysms. Report of four cases. *J Neurosurg.* 2002;97:692-6.

Figure 1:

Panel A: Illustration of the subarachnoid hemorrhage (SAH) distributed mostly around the right cerebello-pontine angle (solid arrow) on native head-CT hours after onset of severe headaches. Panel B: digital-subtraction angiography image in the anterior-posterior plane obtained the day after diagnosing the SAH showing the course of the right anterior inferior cerebellar artery (AICA) (dashed arrow) including the ruptured flow-related meatal AICA-aneurysm (solid arrow) and the arterio-venous malformation (star) fed predominantly by the AICA. Panels C&D: axial and coronal T1 post-contrast MR-images demonstrating the right AICA, the flow-related AICA-aneurysm (solid arrow) and its close proximity to the vestibulo-cochlear nerve (dashed-filled arrows) and the facial nerve (dashed-empty arrow). Panel E: demonstration of the coils (dashed arrow) after successful occlusion of the AICA-aneurysm located within the proximal part of the internal auditory canal.

